

Effective action approach to a bi-layer frustrated antiferromagnet

A. V. Dotsenko

School of Physics, The University of New South Wales, Sydney 2052, Australia
(Internet: `dav@newt.phys.unsw.edu.au`)

Abstract

A bi-layer quantum frustrated antiferromagnet is studied using an effective action approach. The action derived from the microscopical Hamiltonian has the form of the $O(3)$ non-linear sigma model. It is solved in the mean field approximation with the ultraviolet cut off chosen to fit numerical results. The obtained phase diagram displays a decrease in the critical value of inter-layer coupling with increase of in-plane frustration. The critical point for a single-layer frustrated antiferromagnet (the J_1 - J_2 model) is estimated to be $J_{2c} = 0.19J_1$.

PACS Numbers: 75.50.Ee, 75.10.Jm, 75.30.Kz, 75.10.-b

In the past few years, a lot of attention has been given to order-disorder transitions in two-dimensional antiferromagnets. Much of the interest stems from relevance of the problem to high-temperature superconductors where such transitions occur. Although in superconductors the transition is driven by mobile holes, it can also be studied in a frustrated antiferromagnet. The antiferromagnet with frustrating interaction between next nearest neighbours (the J_1 - J_2 model) is the simplest case and has often been studied.¹

In this paper, I study a *bi-layer* frustrated antiferromagnet. The bi-layer problem has emerged from the studies of $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$ which consists of pairs of close CuO_2 planes. Experimentally, there are a number of differences between bi-layer and single layer compounds.² Recent infrared measurements³ suggest that the antiferromagnet interlayer coupling within bi-layers may be as strong as 60 meV (about half of the in-plane coupling) in which case it must be taken into account in any theoretical study. Several authors have discussed pairing in bi-layer models.⁴ This work is restricted to studying magnetic order in the static antiferromagnet.

The Hamiltonian is taken in the form

$$H = J_1 \sum_{\text{NN}} (\mathbf{S}_{1\mathbf{r}} \cdot \mathbf{S}_{1\mathbf{r}'} + \mathbf{S}_{2\mathbf{r}} \cdot \mathbf{S}_{2\mathbf{r}'}) + J_2 \sum_{2\text{N}} (\mathbf{S}_{1\mathbf{r}} \cdot \mathbf{S}_{1\mathbf{r}'} + \mathbf{S}_{2\mathbf{r}} \cdot \mathbf{S}_{2\mathbf{r}'}) \quad (1)$$

$$+ J_3 \sum_{3\text{N}} (\mathbf{S}_{1\mathbf{r}} \cdot \mathbf{S}_{1\mathbf{r}'} + \mathbf{S}_{2\mathbf{r}} \cdot \mathbf{S}_{2\mathbf{r}'}) + J_{\perp} \sum_{\mathbf{r}} \mathbf{S}_{1\mathbf{r}} \cdot \mathbf{S}_{2\mathbf{r}},$$

where the summation is performed over a square lattice, NN denotes nearest neighbours, 2N (3N) stands for second (third) neighbours, and the two planes within the bi-layer are labelled as 1 and 2 (it is assumed that all interactions J are positive).

In case $J_{\perp} = 0$ the Hamiltonian (1) reduces to a frustrated Heisenberg antiferromagnet. Increasing J_2 and J_3 leads to a quantum phase transition into a disordered phase. The numerical value of the critical coupling J_{2c} (at $J_3 = 0$) is, however, known very poorly.

The case of $J_2 = J_3 = 0$ (a bi-layer antiferromagnet) has also received a lot of attention.⁵⁻¹² Availability of reliable numerical results^{8,10} for this case makes it a convenient model¹¹ for testing theoretical results. The bi-layer antiferromagnet has also been used⁹ to explain some experimental results in bi-layer superconductors. However, the value of interlayer coupling required for this ($J_{\perp} \gtrsim 2.5J_1$) is too large to be realistic. Apparently, both in-plane and inter-plane effects are important. In the presence of in-plane frustration the critical value of interlayer coupling may be drastically reduced.

In both the bi-layer antiferromagnet and single layer frustrated antiferromagnet, the mean field spin wave and mean field Schwinger boson theories significantly overestimate the region of stability of the Néel ordered phase. The main reason for this appears to be that these theories assume long range order from the start and do not include some types of fluctuations (see discussion of longitudinal spin fluctuations in bi-layer antiferromagnet in Ref. 12).

In this study, I use an effective action approach which is based on mapping the microscopic Hamiltonian onto the quantum $O(3)$ non-linear sigma model and which has been extensively applied to two-dimensional antiferromagnets. Using a $1/N$ expansion of the non-linear sigma model, Chubukov, Sachdev, and Jinwu Ye¹³ have recently presented a comprehensive analysis of the general properties of clean two-dimensional antiferromagnets

in the vicinity of the order-disorder transition. Good agreement with numerical and experimental data has been obtained.

In this work, I first derive the effective action for the bi-layer frustrated antiferromagnet. It is observed then that for the purposes of the study, the action can be reduced to that of the standard one-band non-linear sigma model. The phase diagram is then found using the mean field (saddle point) solution with the ultraviolet cut off (or, equivalently, the critical coupling) adjusted so as to fit numerical results for the bi-layer antiferromagnet.

There are several ways to obtain the action. I follow the derivation based on coherent state representation.¹⁴ Each spin is represented by a single unit vector $\mathbf{N}$. Then all spin configurations are expressed as a combination of four fields (two fields per layer)

$$\begin{aligned}\mathbf{N}_{1\mathbf{r}} &= \eta_{\mathbf{r}} \mathbf{n}_{1\mathbf{r}} (1 - a^2 \mathbf{L}_{1\mathbf{r}}^2)^{1/2} + a \mathbf{L}_{1\mathbf{r}}, \\ \mathbf{N}_{2\mathbf{r}} &= -\eta_{\mathbf{r}} \mathbf{n}_{2\mathbf{r}} (1 - a^2 \mathbf{L}_{2\mathbf{r}}^2)^{1/2} + a \mathbf{L}_{2\mathbf{r}},\end{aligned}\tag{2}$$

where

$$\mathbf{n}_1^2 = \mathbf{n}_2^2 = 1, \quad \mathbf{n}_1 \cdot \mathbf{L}_1 = \mathbf{n}_2 \cdot \mathbf{L}_2 = 0, \quad a^2 \mathbf{L}_{1,2}^2 \ll 1,$$

$\eta_{\mathbf{r}} = \pm 1$ for the two sublattices, and a is the lattice constant. The unit length fields $\mathbf{n}_{i\mathbf{r}}$ in the classical limit describe orientation of the local Néel ordering and the fields $\mathbf{L}_{i\mathbf{r}}$ represent small local fluctuations.

Now the Hamiltonian is expressed using Eq. (2) and the continuum limit is taken by making an expansion in gradients of $\mathbf{n}_1$ and $\mathbf{n}_2$ and powers of $\mathbf{L}_1$ and $\mathbf{L}_2$

$$\begin{aligned}H &= \int d^2\mathbf{r} \left[\frac{1}{2} \rho_{s1} [(\nabla \mathbf{n}_1)^2 + (\nabla \mathbf{n}_2)^2] + S^2 J_{\perp} (\mathbf{n}_1 - \mathbf{n}_2)^2 \right. \\ &\quad \left. + 4S^2 (J_1 + \frac{1}{8} J_{\perp}) (\mathbf{L}_1^2 + \mathbf{L}_2^2) + S^2 J_{\perp} \mathbf{L}_1 \cdot \mathbf{L}_2 \right],\end{aligned}\tag{3}$$

where $\rho_{s1} = S^2(J_1 - 2J_2 - 4J_3)$ is the bare spin stiffness.

Decomposition of the partition function is a straightforward extension of the procedure for the single layer antiferromagnet¹⁴ and leads to

$$Z = \int \mathcal{D}\mathbf{n}_1 \mathcal{D}\mathbf{n}_2 \mathcal{D}\mathbf{L}_1 \mathcal{D}\mathbf{L}_2 \delta(\mathbf{n}_1^2 - 1) \delta(\mathbf{n}_2^2 - 1) \exp(S_n)\tag{4}$$

with the action

$$S_n = S_B - \int d^2\mathbf{r} \int_0^{\beta} d\tau [H(\tau) + iS\mathbf{L}_1 \cdot (\mathbf{n}_1 \times \partial_{\tau} \mathbf{n}_1) + iS\mathbf{L}_2 \cdot (\mathbf{n}_2 \times \partial_{\tau} \mathbf{n}_2)],$$

where S_B is the residual Berry phase which is zero for all smooth spin configurations and which is ignored hereafter. Finally, the fields $\mathbf{L}_1$ and $\mathbf{L}_2$ are integrated out of Eq. (4)

$$\begin{aligned}Z &= \int \mathcal{D}\mathbf{n}_1 \mathcal{D}\mathbf{n}_2 \delta(\mathbf{n}_1^2 - 1) \delta(\mathbf{n}_2^2 - 1) \\ &\quad \times \exp \left\{ - \int d^2\mathbf{r} \int_0^{\beta} d\tau \left[\frac{1}{2} \rho_{s1} [(\nabla \mathbf{n}_1)^2 + (\nabla \mathbf{n}_2)^2] + S^2 J_{\perp} (\mathbf{n}_1 - \mathbf{n}_2)^2 \right. \right. \\ &\quad \left. \left. + \frac{1}{8} (J_1 + \frac{1}{4} J_{\perp})^{-1} (\mathbf{n}_1 \times \partial_{\tau} \mathbf{n}_1 + \mathbf{n}_2 \times \partial_{\tau} \mathbf{n}_2)^2 + \frac{1}{8} J_1^{-1} (\mathbf{n}_1 \times \partial_{\tau} \mathbf{n}_1 - \mathbf{n}_2 \times \partial_{\tau} \mathbf{n}_2)^2 \right] \right\}.\end{aligned}\tag{5}$$

At $J_{\perp} = 0$, the action (5) represents two independent sigma models each possessing a Goldstone mode at $T = 0$ due to spontaneous symmetry breaking. At finite $J_{\perp}$, one mode acquires a gap $\Delta \propto \sqrt{J_{\perp}J_1}$. There is no general solution in this case because of the presence of the term $(\mathbf{n}_1 \times \partial \mathbf{n}_1) \cdot (\mathbf{n}_2 \times \partial \mathbf{n}_2)^2$ in the action. However, at larger $J_{\perp}$ and $T \ll \sqrt{J_{\perp}J_1}$ the action can be simplified due to the fact that it is dominated by configurations with $\mathbf{n}_1 \approx \mathbf{n}_2$. In this regime of coupled planes, I restrict myself to the Goldstone mode and set $\mathbf{n} = \mathbf{n}_1 = \mathbf{n}_2$ in Eq. (5). The action takes the form

$$Z = \int \mathcal{D}\mathbf{n} \delta(\mathbf{n}^2 - 1) \exp \left\{ -\frac{1}{2g} \int d^2\mathbf{r} \int_0^{c\beta} d\tau [(\nabla \mathbf{n})^2 + (\partial_{\tau} \mathbf{n})^2] \right\}, \quad (6)$$

where time has been rescaled ($c\tau \rightarrow \tau$) and the coupling constant is $g = c/\rho_s$ with the spin wave velocity

$$c = 2S[2(J_1 - 2J_2 - 4J_3)(J_1 + \frac{1}{4}J_{\perp})]^{1/2} \quad (7)$$

and the bare spin stiffness

$$\rho_s = 2S^2(J_1 - 2J_2 - 4J_3). \quad (8)$$

From Eq. (7) for the bi-layer antiferromagnet the spin wave velocity at the critical point $J_{\perp} = 2.5J_1$ is $c_c = 1.8J_1$. It is in good agreement with numerical results¹⁰ which suggests that $1/S$ corrections are negligible at this point. The spin stiffness Eq. (8) is twice as large as in the regime of independent planes. Thus when $J_{\perp}$ is introduced the system first becomes more “classical” (in other words, the effective spin S^* increases) and then evolves towards the quantum disordered phase. This effect was observed in earlier studies^{7,9,12} at a fairly small value of $J_{\perp} \approx 0.1J_1$. With in-plane frustration present the crossover will occur at even smaller values of $J_{\perp}$. (In a recent exact diagonalisation study of the bi-layer t - J model,¹⁵ it was noticed that the two layers become essentially correlated at $J_{\perp} \approx 0.2J$.)

Equation (6) represents the quantum $O(3)$ non-linear sigma model. Not being interested in details of the critical behaviour, I use the simplest mean field solution which is exact for the $O(\infty)$ non-linear sigma model. An ultraviolet cut off must be introduced in momentum integration and a Pauli-Villars cut off is a convenient choice.¹⁴ In the mean field approximation, there are spin-1 excitations with a gapped spectrum and no damping. Straightforward calculations show that the gap m is determined from

$$\log \left(\sinh \frac{m}{2T} \right) = -\frac{2\pi}{T} \left(\frac{1}{g} - \frac{1}{g_c} \right) \quad (9)$$

where $g_c = 4\pi/\Lambda$ is the critical coupling (Λ is the cut off). The system is ordered at $T = 0$ when $g < g_c$. Knowing the critical point at $J_2 = J_3 = 0$ allows to eliminate uncertainty from the cut off. Using $J_{\perp c} = 2.5J$, the cut off is found to be $\Lambda = 1.1\pi$. It is easy to see that the critical line $g = g_c$ has the form

$$J - 2J_2 - 4J_3 = \frac{J + \frac{1}{4}J_{\perp}}{J + \frac{1}{4}J_{\perp c}}. \quad (10)$$

TABLE I. The critical coupling in the J_1 - J_2 model obtained by different methods.

method	J_{2c}
Green functions ^a	0.12
renormalization group ^b	0.15
mean field bond operators ^c	0.19
series expansion ^d	0.33 ± 0.08
present work	0.19
exact diagonalisations ^e	> 0.34

^aRef. 16

^bRef. 17

^cRef. 18

^dRef. 19

^eRef. 20

While Eq. (9) is only the mean field solution and is not quite accurate, Eq. (10) is independent of $1/N$ corrections. The phase diagram is presented in Figs. 2 and 3. Figure 2 is the phase diagram for $J_3 = 0$. In the absence of interlayer coupling, the critical value of J_2 is found to be $J_{2c} = 0.19J_1$. It is compatible with the results obtained by other methods except for numerical results (see Table I). However, numerical studies²⁰ face severe size restrictions and are not very reliable quantitatively. Figure 3 is the phase diagram for $J_2 = 0$. Both Figs. 2 and 3 show that in the presence of an interlayer coupling smaller values of in-plane frustration are sufficient for the disordering transition.

In conclusion, I have studied the bi-layer frustrated antiferromagnet using the effective action approach. It was demonstrated that the essential physics can be described by a one-band non-linear sigma model. The region of stability of the Néel ordered phase has been identified. The critical value of next nearest neighbour interaction for the J_1 - J_2 model is estimated to be $J_2 = 0.19J_1$.

ACKNOWLEDGEMENTS

I am grateful to O. P. Sushkov for discussions and to A. V. Chubukov for his comments on the paper. This work forms part of a project supported by a grant of the Australian Research Council.

REFERENCES

- ¹ A. V. Dotsenko and O. P. Sushkov, Phys. Rev. B **50**, 13 821 (1994) and references therein.
- ² T. Siegrist et al, Phys. Rev. B **35**, 7137 (1989). J. M. Tranquada, G. Shirane, B. Keimer, S. Shamoto, and M. Sato, Phys. Rev. B **40**, 4503 (1989)
- ³ M. Grüninger, J. Münzel, A. Gaymann, A. Zibold, H. P. Geserich, and T. Kopp, preprint MG-94-1, `cond-mat/9501065`.
- ⁴ K. Kuboki and P. A. Lee, `cond-mat/9501030`. A. I. Liechtenstein, I. I. Mazin, and O. K. Andersen, Phys. Rev. Lett. (in press); `cond-mat/9501118`. R. R. dos Santos, `cond-mat/9502043`.
- ⁵ K. Hida, J. Phys. Soc. Jpn, **59**, 2230 (1990).
- ⁶ T. Matsuda and K. Hida, J. Phys. Soc. Jpn **59**, 2223 (1990).
- ⁷ Shi-Min Cui, Commun. in Theor. Phys. **16**, 401 (1991).
- ⁸ K. Hida, J. Phys. Soc. Jpn **61**, 1013 (1992).
- ⁹ A. J. Millis and H. Monien, Phys. Rev. Lett. **70**, 2810 (1993); Phys. Rev. B **50**, 16 606 (1994).
- ¹⁰ A. W. Sandvik and D. J. Scalapino, Phys. Rev. Lett. **72**, 2777 (1994).
- ¹¹ A. W. Sandvik, A. V. Chubukov, and S. Sachdev, `cond-mat/9502012`.
- ¹² A. V. Chubukov and D. K. Morr, `cond-mat/9503029`.
- ¹³ A. V. Chubukov, S. Sachdev, and Jinwu Ye, Phys. Rev. B **49**, 11 919, (1994).
- ¹⁴ S. Sachdev, *Low Dimensional Quantum Field Theories for Condensed Matter Physicists*, Proceedings of the Trieste Summer School 1992 (World Scientific, Singapore, 1993). `cond-mat/9303014`.
- ¹⁵ R. Eder, Y. Ohta, and S. Maekawa, `cond-mat/9504013`.
- ¹⁶ A. F. Barabanov and V. M. Berezovsky, Phys. Lett. A, **186**, 175 (1994); Zhur. Eksp. Teor. Fiz. **106**, 1156 (1994) [JETP **79**, 627 (1994)]; J. Phys. Soc. Jpn **63**, 3974 (1994).
- ¹⁷ T. Einarsson, P. Fröjdh, and H. Johannesson, Phys. Rev. B **45**, 13 121 (1992).
- ¹⁸ S. Sachdev and R. N. Bhatt, Phys. Rev. B **41**, 9323 (1990).
- ¹⁹ M. P. Gelfand, R. R. P. Singh, and D. A. Huse, Phys. Rev. B **40**, 10 801 (1989); J. Stat. Phys. **59**, 1093 (1990); M. P. Gelfand, Phys. Rev. B **42**, 8206 (1990).
- ²⁰ H. J. Schulz, T. A. L. Ziman, and D. Poilblanc, `cond-mat/9402061`.

FIG. 1. A scheme of the bi-layer antiferromagnet.

FIG. 2. The phase diagram for $J_3 = 0$.

FIG. 3. The phase diagram for $J_2 = 0$.

FIGURE 1




FIGURE 2




FIGURE 3

